

STRONGLY MEAGER SETS CAN BE QUITE BIG

TOMEK BARTOSZYNSKI, ANDRZEJ NOWIK, AND TOMASZ WEISS

ABSTRACT. Assume CH. There exists a strongly meager set $X \subseteq 2^\omega$ and a continuous function $F : 2^\omega \rightarrow 2^\omega$ such that $F'''(X) = 2^\omega$.

1. INTRODUCTION

A set $X \subseteq 2^\omega$ is strongly meager ($X \in \mathcal{SM}$) if for every null set $H \in \mathcal{N}$, $X + H \neq \emptyset$. A set $X \subseteq 2^\omega$ is strongly measure zero ($X \in \mathcal{SN}$) if for every meager set $F \in \mathcal{M}$, $X + F \neq \emptyset$.

Let $\mathcal{I} = \{X \subseteq 2^\omega : \forall F : 2^\omega \rightarrow 2^\omega \text{ continuous, } F'''(X) \neq 2^\omega\}$.

The following is well-known:

Lemma 1. 1. $\mathcal{I}$ is a σ -ideal,
 2. $\mathcal{I} \subseteq (s)_0$,
 3. $\mathcal{I} \neq (s)_0$ (in ZFC).

Notice that such a σ -ideal was defined and investigated in several papers, see for example [4].

Since strongly meager sets and strong measure zero sets are $(s)_0$ it makes sense to ask if they are in $\mathcal{I}$.

It is well-known that $\mathcal{SN} \subseteq \mathcal{I}$. In fact, if $F : 2^\omega \rightarrow 2^\omega$ is a continuous function and $X \in \mathcal{SN}$ then $F'''(X) \in \mathcal{SN}$.

The purpose of this paper is to show:

Theorem 2. *It is consistent with ZFC that $\mathcal{SM} \not\subseteq \mathcal{I}$.*

Let us remark that we also have,

Theorem 3 ([2]). *It is consistent with ZFC that $\mathcal{I} = \mathcal{SM}$.*

It is also easy to see that (under CH for example) $\mathcal{I} \not\subseteq \mathcal{SM}$.

2. COMBINATORICS

The following theorem is the finitary version of the construction.

Theorem 4. *For every $k \in \omega$, $\varepsilon, \delta > 0$ there exists $n \in \omega$ such that if $I \subseteq \omega$, $|I| > n$ then there is a partition $2^I = a^0 \dot{\cup} a^1$ such that*

$$1. \text{ if } X \subseteq 2^I, |X| \leq k \text{ then } \left| \frac{|\bigcap_{x \in X} (a^0 + x)|}{2^{|I|}} - \frac{1}{2^{|X|}} \right| < \delta,$$

2000 *Mathematics Subject Classification.* 2000 Mathematics Subject Classification: 03E15, 03E20, 28E15.

Key words and phrases. Strongly meager sets, uniformly continuous images.

The first author was partially supported by NSF grant DMS 9971282 and Alexander von Humboldt Foundation.

2. If $U \subseteq 2^I$, and $\frac{|U|}{2^{|I|}} = a \geq \varepsilon$ then there exists a set $T_U \subseteq 2^I$, $\frac{|T_U|}{2^{|I|}} > 1 - \delta$ such that for every $s \in T_U$ $\left| \frac{|(a^0 + s) \cap U|}{2^{|I|}} - \frac{a}{2} \right| < \delta$.

Proof. (1) This is a special case of a result proved in [3] Fix k, ε, δ , and choose the set $C \subseteq 2^I$ randomly (for the moment I is arbitrary). For each $s \in 2^I$ decisions whether $s \in C$ are made independently with the probability of $s \in C$ equal to $1/2$. Thus the set C is a result of a sequence of Bernoulli trials. Note that by the Chebyshev's inequality, the probability that $1/2 + \delta \geq |C| \cdot 2^{-|I|} \geq 1/2 - \delta$ approaches 1 as $|I|$ goes to infinity.

Let S_n be the number of successes in n independent Bernoulli trials with probability of success p . We will need the following well-known fact.

Theorem 5 (Bernstein's Inequality). *For every $\delta > 0$,*

$$P\left(\left|\frac{S_n}{n} - p\right| \geq \delta\right) \leq 2e^{-n\delta^2/4}.$$

Consider an arbitrary set $X \subseteq 2^I$. To simplify the notation denote $V = 2^I \setminus C$ and note that $\bigcap_{s \in X} (C + s) = 2^I \setminus (V + X)$. For a point $t \in 2^I$, $t \notin X + V$ is equivalent to $(t + X) \cap V = \emptyset$. Thus the probability that $t \notin X + V$ is equal to $2^{-|X|}$, as $t \notin X + V$ means that $t + x \notin V$ for $x \in X$.

Let $G(X)$ be a subgroup of $(2^I, +)$ generated by X . Since every element of 2^I has order 2, it follows that $|G(X)| \leq 2^{|X|}$.

Lemma 6. *There are sets $\{U_j : j \leq |G(X)|\}$ such that:*

1. $\forall j \forall s, t \in U_j \ (s \neq t \rightarrow s + t \notin G(X))$,
2. $\forall j \leq |G(X)| \ |U_j| = 2^{|I|}/|G(X)|$,
3. $\forall i \neq j \ U_i \cap U_j = \emptyset$,
4. $\bigcup_{j \leq |G(X)|} U_j = 2^I$.

PROOF Choose U_j 's to be disjoint selectors from the cosets $2^I/G(X)$. $\square$

Note that if $t_1, t_2 \in U_j$ then the events $t_1 \in X + V$ and $t_2 \in X + V$ are independent since sets $t_1 + X$ and $t_2 + X$ are disjoint. Consider the sets $X_j = U_j \cap \bigcap_{s \in X} (C + s)$ for $j \leq |G(X)|$. The expected value of the size of this set is $2^{-|X|} \cdot 2^{|I|}/|G(X)|$. By theorem 5 for each $j \leq |G(X)|$,

$$P\left(\left|\frac{|X_j|}{2^{|I|}/|G(X)|} - 2^{-|X|}\right| \geq \delta\right) \leq 2e^{-2^{|I|-2}\delta^2/|G(X)|}.$$

It follows that for every $X \subseteq 2^I$ the probability that

$$2^{-|X|} - \delta \leq \frac{|\bigcap_{s \in X} (C + s)|}{2^{|I|}} \leq 2^{-|X|} + \delta$$

is at least

$$1 - 2|G(X)|e^{-2^{|I|-2}\delta^2/|G(X)|} \geq 1 - 2^{|X|+1}e^{-2^{|I|-|X|-2}\delta^2}.$$

The probability that it happens for every X of size $\leq k$ is at least

$$1 - 2^{|I| \cdot (k+1)^2} \cdot e^{-2^{|I|-k-2}\delta^2}.$$

If k and δ are fixed then this expression approaches 1 as $|I|$ goes to infinity, since $\lim_{x \rightarrow \infty} P(x)e^{-x} = 0$ for any polynomial $P(x)$. It follows that for sufficiently large

$|I|$ the probability that the “random” set C has the required properties is > 0 . Thus there exists an actual C with these properties as well.

(2) Let

$$A = \{(s, t) : t \in a^0 + s\}$$

Note that A is symmetric, that $(A)_s = \{t : (s, t) \in A\} = (A)^s = \{t : (t, s) \in A\}$.

Let $U \subseteq 2^I$ be such that $\frac{|U|}{2^{|I|}} = a \geq \varepsilon$. Consider sets $(A)^s = a^0 + s$ for $s \in U$. We want to know how many vertical sections of the set $A \cap (2^I \times U)$ are of size approximately $1/2$ relative to U . Fix $s \in 2^I$ and consider the set $\frac{|U \cap (A)_s|}{|U|}$. By the choice of a_0 the events $u \in a_0 + s$ (equivalent to $u + s \in a_0$) are pairwise independent with probability $1/2$. Thus, by theorem 5,

$$P\left(\left|\frac{|(A)_s \cap U|}{|U|} - \frac{1}{2}\right| \geq \delta\right) < 2e^{-\frac{|U|\delta^2}{4}}.$$

Probability that it holds for some s is at most $2^{|I|+1}e^{-\frac{|U|\delta^2}{4}}$. Since $\frac{|U|}{2^{|I|}} = a \geq \varepsilon$ we get

$$\begin{aligned} P\left(\left|\frac{|(A)_s \cap U|}{|U|} - \frac{1}{2}\right| \geq \delta\right) &= P\left(\left|\frac{|(A)_s \cap U|}{2^{|I|}} \cdot \frac{1}{a} - \frac{1}{2}\right| \geq \delta\right) = \\ &P\left(\left|\frac{|(A)_s \cap U|}{2^{|I|}} - \frac{a}{2}\right| \geq a\delta\right) \leq 2^{|I|+1}e^{-\frac{|U|\delta^2}{4}} \leq 2^{|I|+1}e^{-\frac{2^{|I|}\varepsilon\delta^2}{4}} \end{aligned}$$

It follows that

$$P\left(\left|\frac{|(A)_s \cap U|}{2^{|I|}} - \frac{a}{2}\right| \geq \delta\right) \leq 2^{|I|+1}e^{-\frac{2^{|I|}\varepsilon\delta^2}{4}} \rightarrow 0 \text{ as } |I| \rightarrow \infty.$$

□

Definition 7 ([3]). Suppose that I is a finite set. A distribution is a function $m : I \rightarrow \mathbb{R}$ such that

$$0 \leq m(x) \leq \frac{1}{|I|}.$$

Let $\bar{m} = \sum_{s \in I} m(s)$.

To illustrate this concept suppose that $A \subseteq 2^\omega$ is a measurable set and $n \in \omega$. Define m on 2^n by $m(s) = \mu(A \cap [s])$ for $s \in 2^n$. A specific instance of this definition that we will use often in the sequel is when A is a clopen set. In particular, if $K = I_1 \cup I_2$ and $J \subseteq 2^K$, define distribution m on 2^{I_1} as follows: for $s \in 2^{I_1}$ let

$$m(s) = \frac{|\{t \in J : s \subseteq t\}|}{2^{|K|}}.$$

The following theorem is an extension of theorem 4 dealing with distributions instead of sets.

Theorem 8. For every $k \in \omega$, $\varepsilon, \delta > 0$ there exists $n \in \omega$ such that if $I \subseteq \omega$, $|I| > n$ then there is a partition $2^I = a^0 \cup a^1$ such that

1. if $X \subseteq 2^I$, $|X| \leq k$ then $\left|\frac{|\bigcap_{x \in X}(a^0 + x)|}{2^{|I|}} - \frac{1}{2^{|X|}}\right| < \delta$,

2. If m is a distribution on 2^I , and $\overline{m} \geq \varepsilon$ then there exists a set $T_m \subseteq 2^I$, $\frac{|T_m|}{2^{|I|}} > 1 - \delta$ such that for every $s \in T_m$, $\left| \sum \{m(t) : t \in a^0 + s\} - \frac{\overline{m}}{2} \right| < \delta$.

Proof. This is a generalization of theorem 4. Suppose that k, ε, δ are given and $a_0 \dot{\cup} a_1 = 2^I$ are as in theorem 4. First observe that if $m = \frac{b}{2^{|I|}} \cdot \chi_U$, where $U \subseteq 2^I$ and $\frac{|U|}{2^{|I|}} = a \geq \varepsilon$, $0 < b \leq 1$ and χ_U is a characteristic function of the set U then it follows immediately from theorem 4 that for $s \in T_U$,

$$\left| \sum \{m(t) : t \in a^0 + s\} - \frac{\overline{m}}{2} \right| < b\delta \leq \delta.$$

Next, note that if $\{U_i : i < \ell\}$, $\{b_i : i < \ell\}$ are such that

1. $U_i \cap U_j = \emptyset$ for $i \neq j$,
2. $0 < b_i \leq 1$ for $i < \ell$,
3. $\frac{|U_i|}{2^{|I|}} \geq \varepsilon$ for every $i < \ell$,
4. $m_i = \frac{b_i}{2^{|I|}} \cdot \chi_{U_i}$.

then for $m = \sum_{i < \ell} m_i$ and $s \in \bigcap_{i < \ell} T_{U_i}$ we have

$$\left| \sum \{m(t) : t \in a^0 + s\} - \frac{\overline{m}}{2} \right| < \sum_{i < \ell} b_i \delta \leq \ell \cdot \delta \quad \text{and} \quad \frac{|\bigcap_{i < \ell} T_{U_i}|}{2^{|I|}} \geq 1 - \ell \delta.$$

Fix $k \in \omega, \varepsilon > \delta > 0$. Apply theorem 4 for k , $\varepsilon' = \delta^2$ and $\delta' = \delta^3$ to get n and $a_0 \dot{\cup} a_1 = 2^I$. We will show that a_0, a_1 satisfy the requirements of the theorem.

Consider an arbitrary distribution m and let $\ell = \frac{1}{\delta}$ (without loss of generality it is an integer). Let $\{U_i : i < \ell\}$ be defined as

$$U_i = \left\{ s \in 2^I : \frac{i\delta}{2^{|I|}} < m(s) \leq \frac{(i+1)\delta}{2^{|I|}} \right\}.$$

Let $K = \left\{ i : \frac{|U_i|}{2^{|I|}} \geq \delta^2 \right\}$. Put $m' = \sum_{i \in K} \frac{i\delta}{2^{|I|}} \cdot \chi_{U_i}$ and $U = \bigcup_{i < \ell} U_i$. Note that for $s \in U$, and $i \in K$, $m(s) - m'(s) \leq \frac{\delta}{2^{|I|}}$ and $|\overline{m} - \overline{m'}| \leq \delta + \sum_{i < \ell} \delta^2 = 2\delta$.

Apply 4 to each of the sets $\{U_i : i \in K\}$ to get sets T_{U_i} and put $T_m = \bigcap_{i \in K} T_{U_i}$. Clearly $\frac{|T_m|}{2^{|I|}} \geq 1 - \ell \cdot \delta^3 = 1 - \delta^2$. Now, for $t \in T_m$,

$$\sum \{m(s) : s \in a^0 + t\} \leq \sum \{m'(s) : s \in a^0 + t\} + 2\delta \leq \frac{\overline{m'}}{2} + \delta^3 + 2\delta + \delta^2 \leq \frac{\overline{m}}{2} + 3\delta.$$

Lower estimate is the same and we get

$$\left| \sum \{m(s) : s \in a^0 + t\} - \frac{\overline{m}}{2} \right| \leq 3\delta.$$

□

3. ZFC RESULT

As a warm-up before proving the main result we will show a ZFC result that uses only small portion of the combinatorial tools developed above.

In order to show that $\mathcal{SN} \subseteq \mathcal{I}$ one could use the following result:

Theorem 9 ([5]). *Suppose that $F : 2^\omega \longrightarrow 2^\omega$ is a continuous function. There exists a set $H \in \mathcal{M}$ such that*

$$\forall z \in 2^\omega \exists y \in 2^\omega F^{-1}(y) \subseteq H + z.$$

We will show that the measure analog of this theorem is false.

Theorem 10. *There exists a continuous function $F : 2^\omega \longrightarrow 2^\omega$ such that for every set $G \in \mathcal{N}$,*

$$\{z : \exists y F^{-1}(y) \subseteq G + z\} \in \mathcal{N}.$$

Proof. Let δ_n, k_n, I_n for $n \in \omega$ be such that

1. $\delta_n = 4^{-n-3}$, $k_n = 2^{n+3}$,
2. I_n is chosen as in Theorem 4(1) for $k = k_n$ and $\delta = \delta_n$ and
 - (a) $|I_{n+1}| \geq 4^{n+10} 2^{|I_0 \cup \dots \cup I_n|}$
 - (b) $\min(I_{n+1}) \geq \max(I_n)$.

Let $2^{I_n} = a_n^0 \dot{\cup} a_n^1$ be a partition as in theorem 4(1).

Define $F : 2^\omega \longrightarrow 2^\omega$ as

$$F(x)(n) = i \iff x \upharpoonright I_n \in a_n^i.$$

Note that for every $x \in 2^\omega$, $F^{-1}(x) = \prod_n a_n^{x(n)}$ is a perfect set.

Suppose that $G \subseteq 2^\omega$ is null, and let $U \subseteq 2^\omega$ be an open set of measure $1/2 > \varepsilon > 0$ containing G . We will show that

$$\mu(\{z : \exists y F^{-1}(y) \subseteq U + z\}) \longrightarrow 0 \text{ as } \varepsilon \rightarrow 0.$$

Let

$$U_0^\emptyset = \left\{ s \in 2^{I_0} : \frac{\mu([s] \cap U)}{\mu([s])} \geq \frac{3}{4} \right\},$$

and for $n > 0$ and $t \in 2^{I_0 \cup \dots \cup I_{n-1}}$ let

$$U_n^t = \left\{ s \in 2^{I_n} : \frac{\mu([t \frown s] \cap U)}{\mu([t \frown s])} \geq 1 - \frac{1}{2^{n+2}} \right\}.$$

Easy computation shows that $\mu(U_0^\emptyset) \leq 4/3\varepsilon \leq 2/3$. For $t \in 2^{I_0 \cup \dots \cup I_{n-1}}$ we say that t is good if $t \upharpoonright I_0 \notin U_0^\emptyset$, and for $j \leq n-1$, $t \upharpoonright I_j \notin U_j^{t \upharpoonright I_0 \cup \dots \cup I_{j-1}}$. In particular, if $t \in 2^{I_0 \cup \dots \cup I_{n-1}}$ is good then by induction we show that

$$\mu(U_n^t) \leq \frac{2^{n+2}}{2^{n+2}-1} \cdot \frac{\mu(U \cap [t])}{\mu([t])} \leq \frac{2^{n+2}}{2^{n+2}-1} \cdot \left(1 - \frac{1}{2^{n+1}}\right) \leq \frac{2^{n+2}-2}{2^{n+2}-1}.$$

For a good sequence $t \in 2^{I_0 \cup \dots \cup I_{n-1}}$ let

$$Z_n^t = \{v \in 2^{I_n} : \exists i \in 2 \ a_n^i + v \subseteq U_n^t\}.$$

By theorem 4(1), for $i = 0, 1$ and $X \subseteq 2^{I_n}$ of size $n+10$,

$$\frac{|X + a_n^i|}{2^{|I_n|}} = 1 - \frac{|\bigcap_{x \in X} (a_n^i + x)|}{2^{|I_n|}} \geq 1 - \frac{1}{2^{n+10}} - \frac{1}{4^{n+3}} \geq \frac{2^{n+2}-2}{2^{n+2}-1}.$$

Therefore, $|Z_n^t| \leq n + 10$. Let $Z_n = \bigcup \{Z_n^t : t \in 2^{I_0 \cup \dots \cup I_{n-1}} \text{ is good}\}$. We get that $|Z_n| \leq (n + 10) \cdot 2^{|I_0 \cup \dots \cup I_{n-1}|}$, so in particular, $|Z_n| \cdot 2^{-|I_n|} \leq 2^{-n}$. Let

$$W_U = \{z \in 2^\omega : \exists n \ z \restriction I_n \in Z_n\}.$$

Note that if $\varepsilon < 2^{-n-|I_0 \cup \dots \cup I_n|}$ then $Z_0 = Z_1 = \dots = Z_n = \emptyset$. Therefore, $\mu(W_U) \rightarrow 0$ as $\varepsilon \rightarrow 0$. The following lemma finishes the proof.

Lemma 11. *If $F^{-1}(y) \subseteq U + z$ then $z \in W_U$.*

Proof. Suppose not. By induction build a branch $r \in z + F^{-1}(y)$ such that $r + z \restriction I_0 \notin (U)_0^\emptyset$, $r + z \restriction I_1 \notin (U)_1^{r+z \restriction I_0}$, etc. Since U is open, it means that $r + z \notin U$. $\square$

$\square$

4. MAIN RESULT

Theorem 12. *Assume that $\text{cov}(\mathcal{N}) = 2^{\aleph_0}$. There exists a set $X \in \mathcal{SM}$ and a continuous function $F : 2^\omega \rightarrow 2^\omega$ such that $F''(X) = 2^\omega$.*

Suppose that we have sequences $\{\varepsilon_n, \delta_n : n \in \omega\}$ and a partition $\{I_n : n \in \omega\}$ such that

1. $2^{|I_0 \cup I_1 \cup \dots \cup I_n|} \cdot \delta_{n+1} < \varepsilon_{n+1} < 2^{-n}$,
2. $2^{|I_0 \cup I_1 \cup \dots \cup I_n|} \cdot \varepsilon_{n+1} < \delta_n$,
3. $I_n, a_n^0 \dot{\cup} a_n^1 = 2^{I_n}$ is chosen for δ_n, ε_n as in theorem 8.

As in theorem 10 define F as $F(x)(n) = i \iff x \restriction I_n \in a_n^i$.

Lemma 13. *Suppose that a σ -ideal $\mathcal{I}$ on 2^ω , and σ -ideals $\mathcal{J}_x$ of $F^{-1}(x)$ are such that the following holds:*

1. For every $G \in \mathcal{N}$, $\{z \in 2^\omega : \exists x (G + z) \cap F^{-1}(x) \notin \mathcal{J}_x\} \in \mathcal{I}$.
2. for every $G \in \mathcal{N}$ and $t \in 2^\omega$, $\{z : t \in (G + z)\} \in \mathcal{I}$.
3. $\forall x \text{ cov}(\mathcal{J}_x) = \text{cov}(\mathcal{I}) = 2^{\aleph_0}$.

Then there exists a set $X \in \mathcal{SM}$ such that $F''(X) = 2^\omega$.

Proof. Let $\{G_\alpha : \alpha < c\}$ be enumeration of null sets, and $\{t_\alpha : \alpha < c\}$ enumeration of 2^ω . Build by induction sequences $\{x_\alpha, z_\alpha : \alpha < c\}$ such that

1. $x_\alpha \in F^{-1}(t_\alpha)$,
2. $\forall \beta < c \ x_\beta \notin G_\alpha + z_\alpha$.

Suppose that $\{x_\beta, z_\beta : \beta < \alpha\}$ are given.

Consider sets $Z = \{z \in 2^\omega : \exists x (G_\alpha + z) \cap F^{-1}(x) \notin \mathcal{J}_x\}$, and for $\beta < \alpha$, $Z_\beta = \{z : x_\beta \in (G_\alpha + z)\}$. Let $z_\alpha \notin Z \cup \bigcup_{\beta < \alpha} Z_\beta$. Next consider $F^{-1}(t_\alpha)$ and choose $x_\alpha \in F^{-1}(t_\alpha) \setminus \bigcup_{\beta < \alpha} (G_\beta + z_\beta)$. $\square$

In our case $\mathcal{I} = \mathcal{N}$ and $\mathcal{J}_x$ is the measure ideal on $F^{-1}(x)$. In other words, let $c_n = 2^{|I_n|-1}$, and let $\mathcal{J}$ be a σ -ideal of null sets (with respect to the standard product measure) on $\mathcal{X} = \prod_{n=0}^\infty c_n$. Note that $\mathcal{X}$ is chosen to be isomorphic (level by level) with $F^{-1}(x)$ for any x . Let $\mathcal{J}_x$ be the copy of $\mathcal{J}$ on $F^{-1}(x)$.

Specifically, define measure μ_x on $F^{-1}(x)$ as $\mu_x = \prod_{n \in \omega} \mu_n^{x(n)}$, where μ_n^i is a normalized counting measure on a_n^i ($i = 0, 1$). Clearly, μ_x is essentially the Lebesgue measure on $F^{-1}(x)$.

Now what we want to show is that

Lemma 14. *For every $G \in \mathcal{N}$,*

$$\{z : \exists x \mu_x(F^{-1}(x) \cap (G + z)) > 0\} \in \mathcal{N}.$$

Before we go further let us briefly look at the nature of the difficulties in proving the result using theorem 10. The problem is that the relation $F^{-1}(y) \not\subseteq G + z$ is not additive. Quick analysis shows that every choice of a point z that will shift G away from the set we are constructing has to fulfill continuum many requirements. That is why we shift from $F^{-1}(y) \not\subseteq G + z$ to $F^{-1}(y) \cap (G + z) \in \mathcal{J}_y$, an additive requirement. We still have continuum many constraints, this time we need to find z such that $F^{-1}(y) \cap (G + z) \in \mathcal{J}_y$ for every y .

5. PROOF OF LEMMA 14

Suppose that $G \subseteq 2^\omega$ is a null set.

Lemma 15. *There are sequences $\{K_n, K'_n : n \in \omega\}, \{J_n, J'_n : n \in \omega\}$ such that*

1. K_n 's and K'_n 's are consecutive intervals that are unions of I_m 's,
2. $J_n \subseteq 2^{K_n}, J'_n \subseteq 2^{K'_n}$,
3. $\frac{|J_n|}{2^{|K_n|}}, \frac{|J'_n|}{2^{|K'_n|}} < \frac{1}{2^n}$,
4. $G \subseteq H_1 \cup H_2$, where $H_1 = \{x : \exists^\infty n x \restriction K_n \in J_n\}$ and $H_2 = \{x : \exists^\infty n x \restriction K'_n \in J'_n\}$.

Proof. Use the theorem (and its proof) 2.5.7 of [1]. □

Clearly, if we show the lemma 14 for H_1 and for H_2 then we show it for G . Therefore, without loss of generality we can assume that $G = \{x : \exists^\infty n x \restriction K_n \in J_n\}$, where K_n, J_n are as above. Moreover, we can assume that $\frac{|J_n|}{2^{|K_n|}} = \frac{1}{2^n}$, since the property we are interested in reflects downwards.

Now suppose that for $n \in \omega$, $K_n = I_{k_n} \cup \dots \cup I_{k_{n+1}-1}$. Fix $z, x \in 2^\omega$ and let for $n \in \omega$,

$$J_n^{x,z} = (J_n + z \restriction K_n) \cap \prod_{j=k_n}^{k_{n+1}-1} a_j^{x(j)}.$$

Of course $J_n^{x,z}$ depends only on $z \restriction K_n$ and $x \restriction [k_n, k_{n+1})$.

It is easy to see that $F^{-1}(x) \cap (G + z) = \{v \in F^{-1}(x) : \exists^\infty n v \restriction K_n \in J_n^{x,z}\}$. Since $F^{-1}(x) = \prod_n \prod_{j=k_n}^{k_{n+1}-1} a_j^{x(j)}$ it follows that

$$\mu_x(F^{-1}(x) \cap (G + z)) = 0 \iff \sum_n \frac{|J_n^{x,z}|}{\left| \prod_{j=k_n}^{k_{n+1}-1} a_j^{x(j)} \right|} < \infty.$$

Thus we need to find sets $T_n \subseteq 2^{K_n}$ such that $\mu(\prod_n T_n) > 0$ and such that if $z \in \mathbb{Q} + \prod_n T_n$ then

$$\forall x \in 2^\omega \sum_n \frac{|J_n^{x,z}|}{\left| \prod_{j=k_n}^{k_{n+1}-1} a_j^{x(j)} \right|} < \infty.$$

Fix $n \in \omega$ and let $K_n = K = I_k \cup I_{k+1} \cup \dots \cup I_{k+n}$, $J \subseteq 2^K$, $\frac{|J|}{2^{|K|}} = 2^{-n} \geq \varepsilon_k$.

It suffices to show that there exists a set $T_J \subseteq 2^K$ such that $\frac{|T_J|}{2^{|K|}} > 1 - \delta_{k-1}$ and

for every $s \in T_J$, for every $t \in 2^{[k, k+n]}$,

$$\left| \frac{|(J+s) \cap \prod_{j=0}^n a_{k+j}^{t(j)}|}{|\prod_{j=0}^n a_{k+j}^{t(j)}|} - \frac{|J|}{2^{|K|}} \right| < \delta_{k-1}.$$

In this way for $z \in \mathbb{Q} + \prod_n T_n$, and $x \in 2^\omega$, and sufficiently large n ,

$$\left| \frac{|J_n^{x,z}|}{|\prod_{j=k_n}^{k_{n+1}-1} a_j^{x(j)}|} = \frac{|(J+z) \cap \prod_{j=0}^n a_{k+j}^{x(j)}|}{|\prod_{j=0}^n a_{k+j}^{x(j)}|} \leq \frac{|J|}{2^{|K|}} + \delta_{k-1}.$$

It follows that to finish the construction it suffices to prove the lemma below.

Lemma 16. *Suppose that $K = I_k \cup I_{k+1} \cup \dots \cup I_{k+n}$, $J \subseteq 2^K$, $\frac{|J|}{2^{|K|}} \geq \varepsilon_k$. Then there exists a set $T_J \subseteq 2^K$ such that $\frac{|T_J|}{2^{|K|}} > 1 - \delta_{k-1}$ and for every $s \in T_J$,*

$$\left| \frac{|(J+s) \cap \prod_{j=0}^n a_{k+j}^0|}{|\prod_{j=0}^n a_{k+j}^0|} - \frac{|J|}{2^{|K|}} \right| < \delta_{k-1}.$$

Moreover, for every $s \in T_J$, and every $t \in 2^{[k, k+n]}$,

$$\left| \frac{|(J+s) \cap \prod_{j=0}^n a_{k+j}^{t(j)}|}{|\prod_{j=0}^n a_{k+j}^{t(j)}|} - \frac{|J|}{2^{|K|}} \right| < \delta_{k-1}.$$

Proof. For $0 \leq i \leq n$ define distribution m_i on $2^{I_k \cup I_{k+1} \cup \dots \cup I_{k+i}}$ as

$$m_i(s) = \frac{|\{t \in J : s \subseteq t\}|}{2^{|K|}}.$$

Note that $\overline{m_i} = \frac{|J|}{2^{|K|}}$. Observe that the distribution m_n coincides with J , that is,

$$m_n(s) = \begin{cases} \frac{1}{2^{|K|}} & \text{if } s \in J \\ 0 & \text{otherwise} \end{cases}.$$

We will show by induction that for $i \leq n$, there exists a set $T_{m_i} \subseteq 2^{I_k \cup I_{k+1} \cup \dots \cup I_{k+i}}$, $\frac{|T_{m_i}|}{2^{|I_k \cup I_{k+1} \cup \dots \cup I_{k+i}|}} > (1 - \delta_k) \cdot \prod_{j < i} (1 - \delta_{k+j}) > 1 - \delta_{k-1}$ such that for every $s \in T_{m_i}$,

$$\left| \sum \left\{ m_i(t) : t \in \prod_{j=0}^i (a_{k+j}^0 + s \upharpoonright I_{k+j}) \right\} - \frac{1}{2^{i+1}} \cdot \frac{|J|}{2^{|K|}} \right| < \frac{\delta_{k-1}}{2^i}.$$

In particular, for $i = n$, and $s \in T_{m_n} = T_J$,

$$\left| \sum \left\{ m_n(t) : t \in \prod_{j=0}^n (a_{k+j}^0 + s \upharpoonright I_j) \right\} - \frac{1}{2^{n+1}} \cdot \frac{|J|}{2^{|K|}} \right| < \frac{\delta_{k-1}}{2^n}.$$

The last equation means that for $s \in T_J$,

$$\left| \frac{|J \cap (\prod_{j=0}^n a_{k+j}^0 + s \upharpoonright I_J)|}{2^{|K|}} - \frac{1}{2^{n+1}} \cdot \frac{|J|}{2^{|K|}} \right| < \frac{\delta_{k-1}}{2^n}.$$

By moving s , and multiplying by 2^n we finally get,

$$\left| \frac{|(J+s) \cap \prod_{j=0}^n a_{k+j}^0|}{|\prod_{j=0}^n a_{k+j}^0|} - \frac{|J|}{2^{|K|}} \right| < \delta_{k-1}.$$

Let

$$\left| \frac{|(J+s) \cap \prod_{j=0}^n a_{k+j}^0|}{|\prod_{j=0}^i a_{k+j}^0|} - \frac{|J|}{2^{|K|}} \right| = \text{error}_n.$$

We want to show that $\text{error}_n < \delta_{k-1}$.

Before we start induction note that we can shrink J slightly so that the resulting set has the following property,

$$\forall i \leq n \quad \forall s \in 2^{I_k \cup \dots \cup I_{k+i}} \quad \left(m_i(s) \neq 0 \rightarrow m_i(s) \geq \frac{\varepsilon_{k+i}}{2^{|I_k \cup \dots \cup I_{k+i}|}} \right).$$

By removing from J all nodes (and their descendants) that do not have this property we drop the “measure” by $\varepsilon_{k+n} + \varepsilon_{k+n-1} + \dots + \varepsilon_k < 2\varepsilon_k < \frac{\delta_{k-1}}{2}$. So let assume that J has the above property and later add $\frac{\delta_{k-1}}{2}$ to the error term.

The inductive proof is straightforward – for m_0 we get T_{m_0} immediately from theorem 8.

Now consider m_{i+1} . For each $t \in 2^{I_k \cup \dots \cup I_{k+i}}$ let m_{i+1}^t be the distribution on $2^{I_{k+i+1}}$ defined as $m_{i+1}^t(s) = 2^{|I_k \cup \dots \cup I_{k+i}|} m_{i+1}(t \frown s)$.

Clearly, $m_{i+1}^t(s) \leq 2^{|I_k \cup \dots \cup I_{k+i}|} \frac{1}{2^{|I_k \cup \dots \cup I_{k+i+1}|}} = \frac{1}{2^{|I_{k+i+1}|}}$ for every s . Moreover, $\overline{m_{i+1}^t} = 2^{|I_k \cup \dots \cup I_{k+i}|} m_i(t)$. In particular, shrinking J as above yields, if $\overline{m_{i+1}^t} > 0$ then $\overline{m_{i+1}^t} \geq \varepsilon_{k+i}$. For every $t \in 2^{I_k \cup \dots \cup I_{k+i}}$, $\overline{m_{i+1}^t} > 0$ apply theorem 8 to get a set $T_t \subseteq 2^{I_{k+i+1}}$ such that $\frac{|T_t|}{2^{|I_{k+i+1}|}} \geq 1 - \delta_{k+i+1}$, and for every $s \in T_t$,

$$\left| \sum \{m_{i+1}^t(v) : v \in a_{k+i+1}^0 + s\} - \frac{\overline{m_{i+1}^t}}{2} \right| < \delta_{k+i+1}.$$

Let $T_{m_{i+1}} = T_{m_i} \times \bigcap \{T_t : \overline{m_{i+1}^t} > 0\}$. Clearly,

$$\begin{aligned} \frac{|T_{m_{i+1}}|}{2^{|I_k \cup \dots \cup I_{k+i+1}|}} &= \frac{|T_{m_i}|}{2^{|I_k \cup \dots \cup I_{k+i}|}} \cdot \frac{|\bigcap_t T_t|}{2^{|I_{k+i+1}|}} \geq \\ &\left((1 - \delta_k) \cdot \prod_{j < i} (1 - \delta_{k+j}) \right) \cdot \left(1 - 2^{|I_k \cup \dots \cup I_{k+i+1}|} \delta_{k+i+1} \right) \geq \\ &(1 - \delta_k) \cdot \prod_{j \leq i} (1 - \delta_{k+j}) > 1 - \delta_{k-1}. \end{aligned}$$

Suppose that $s \in T_{m_{i+1}}$.

$$\begin{aligned}
& \sum \left\{ m_{i+1}(v) : v \in \prod_{j=0}^{i+1} (a_{k+j}^0 + s \upharpoonright I_{k+j}) \right\} = \\
& \sum \left\{ \sum \{ m_{i+1}(t \cap v) : v \in a_{k+i+1}^0 + s \upharpoonright I_{k+i+1} \} : t \in \prod_{j=0}^i (a_{k+j}^0 + s \upharpoonright I_{k+j}) \right\} = \\
& \sum \left\{ \sum \left\{ \frac{1}{2^{|I_k \cup \dots \cup I_{k+i}|}} m_{i+1}^t(v) : v \in \prod_{j=0}^{i+1} (a_{k+j}^0 + s \upharpoonright I_{k+j}) \right\} : t \in \prod_{j=0}^i (a_{k+j}^0 + s \upharpoonright I_{k+j}) \right\} \leq \\
& \sum \left\{ \frac{1}{2^{|I_k \cup \dots \cup I_{k+i}|}} \frac{\overline{m_{i+1}^t}}{2} + \delta_{k+i+1} : t \in \prod_{j=0}^i (a_{k+j}^0 + s \upharpoonright I_{k+j}) \right\} \leq \\
& \frac{1}{2} \sum \left\{ \frac{1}{2^{|I_k \cup \dots \cup I_{k+i}|}} \overline{m_{i+1}^t} + 2\delta_{k+i+1} : t \in \prod_{j=0}^i (a_{k+j}^0 + s \upharpoonright I_{k+j}) \right\} = \\
& \frac{1}{2} \sum \left\{ m_i(t) + 2\delta_{k+i+1} : t \in \prod_{j=0}^i (a_{k+j}^0 + s \upharpoonright I_{k+j}) \right\} \leq \\
& 2^{|I_k \cup \dots \cup I_{k+i}|} \delta_{k+i+1} + \frac{1}{2} \left(\frac{1}{2^i} \frac{|J|}{2^{|K|}} + \text{error}_i \right) \leq \frac{1}{2^{i+1}} \frac{|J|}{2^{|K|}} + \frac{\text{error}_i}{2} + 2^{|I_k \cup \dots \cup I_{k+i}|} \delta_{k+i+1},
\end{aligned}$$

where error_i is the error term given by the inductive hypothesis. That gives us

$$\text{error}_i \leq \frac{\delta_k}{2^i} + \frac{\delta_{k+1}}{2^{i-1}} + \dots + \delta_{k+i} \leq \frac{\delta_{k-1}}{2^{i+1}},$$

so

$$\sum \left\{ m_{i+1}(v) : v \in \prod_{j=0}^{i+1} (a_{k+j}^0 + s \upharpoonright I_{k+j}) \right\} \leq \frac{1}{2^{i+1}} \frac{|J|}{2^{|K|}} + \frac{\delta_{k-1}}{2^{i+2}}.$$

The lower bound is similar, and we get for $s \in T_{m_{i+1}}$,

$$\left| \sum \left\{ m_{i+1}(t) : t \in \prod_{j=0}^{i+1} (a_{k+j}^0 + s \upharpoonright I_{k+j}) \right\} - \frac{1}{2^{i+1}} \cdot \frac{|J|}{2^{|K|}} \right| < \frac{\delta_{k-1}}{2^{i+2}}.$$

As before that yields the estimate

$$\left| \frac{|(J+s) \cap \prod_{j=0}^n a_{k+j}^0|}{|\prod_{j=0}^n a_{k+j}^0|} - \frac{|J|}{2^{|K|}} \right| < \frac{\delta_{k-1}}{2}.$$

Since we started by reducing the “measure” of J by $\frac{\delta_{k-1}}{2}$ we get the required estimate.

Finally we will show the second part of the lemma. We will proceed by induction on n . If $n = 0$ then $K = I_k$ and by the part already proved

$$\left| \frac{|(J+s) \cap a_k^0|}{|a_k^0|} - \frac{|J|}{2^{|K|}} \right| < \delta_{k-1}.$$

Now

$$\begin{aligned} |(J + s) \cap a_k^1| &= |(J + s) \setminus ((J + s) \cap a_k^0)| \leq |J| - \left(\frac{|J|}{2^{|K|}} - \delta_{k-1} \right) \cdot |a_k^0| \leq \\ &|J + s| - \frac{1}{2}|J| + \delta_{k-1}|a_k^0| = \frac{1}{2}|J| + \delta_{k-1}|a_k^0| \end{aligned}$$

Thus

$$\frac{|(J + s) \cap a_k^1|}{2^{|K|}} \leq \frac{\frac{1}{2}|J| + \delta_{k-1}|a_k^0|}{2^{|K|}} = \frac{1}{2} \frac{|J|}{2^{|K|}} + \frac{\delta_{k-1}}{2}.$$

The lower estimate is similar so we have

$$\left| \frac{|(J + s) \cap a_k^1|}{2^{|K|}} - \frac{1}{2} \frac{|J|}{2^{|K|}} \right| < \frac{\delta_{k-1}}{2},$$

and

$$\left| \frac{|(J + s) \cap a_k^1|}{|a_k^1|} - \frac{|J|}{2^{|K|}} \right| < \delta_{k-1}.$$

The rest of the proof is the repetition of the above argument, the single step computed here shows that there is no difference whether we use a_j^0 or a_j^1 , the estimates do not change. $\square$

REFERENCES

- [1] Tomek Bartoszyński and Haim Judah. *Set Theory: on the structure of the real line*. A.K. Peters, 1995.
- [2] Tomek Bartoszyński and Saharon Shelah. On Borel and dual Borel Conjecture. in preparation.
- [3] Tomek Bartoszyński and Saharon Shelah. Strongly meager sets are not an ideal. *Journal of Mathematical Logic*, 1:1–34, 2001.
- [4] Paul Corazza. The Generalized Borel Conjecture and strongly proper orders. *Transactions of The American Mathematical Society*, 316(1):115–140, 1989.
- [5] Andrzej Nowik and Tomasz Weiss. Strongly meager sets and their uniformly continuous images. *Proceedings of the American Mathematical Society*, 129(1):265–270, 2000.

DEPARTMENT OF MATHEMATICS, BOISE STATE UNIVERSITY, BOISE, IDAHO 83725 U.S.A.

E-mail address: `tomek@math.boisestate.edu`, `http://math.boisestate.edu/~tomek`

Current address: Institute of Mathematics, Polish Academy of Sciences, Abrahamowa 18, 81 – 825 Sopot, Poland

UNIVERSITY OF GDAŃSK, INSTITUTE OF MATHEMATICS, UL. WITA STWOSZA 57, 80 – 952 GDAŃSK, POLAND

E-mail address: `matan@julia.univ.gda.pl`

DEPARTMENT OF MATHEMATICS, WSRP, SIEDLCE

E-mail address: `weiss@wsrp.siedlce.pl`